

A) Analytes	Reproducibility	Direction of FC	Correlation >0.8	Readout shift >40%	Distance to LDD >20 fold
Cortisol	+++++	+	-	X	-
Haptoglobin	+++++	+	+	+	+
Interleukin 10	+++++	+	+	X	X
α -1 antitrypsin	++++	+	+	+	+
Apolipoprotein H	++++	+	X	+	+
CA	++++	+	+	+	X
complement 3	++++	+	X	+	+
Ferritin	++++	+	+	+	+
Interleukin 7	++++	(3/4)	X	+	X
Trail R3	++++	(3/4)	-	+	-
Betacellulin	+++	(2/3)	-	X	-
CTGF	+++	+	-	+	-
Endothelin 1	+++	+	-	X	X
ICAM 1	+++	+	+	+	+
MDC	+++	+	+	+	+
MIF	+++	+	+	+	X
Prolactin	+++	+	+	+	+
Serum Amyloid P	+++	+	+	+	+
Sortilin	+++	(2/3)	+	+	+
TIMP 1	+++	+	+	+	+
TNF R 2	+++	+	+	+	+
VEGF	+++	(2/3)	+	+	+

B) Analytes

Apolipoprotein A1
 BDNF
 Beta-2 Microglobulin
 Fetuin A
 IgA
 LH (Luteinizing Hormone)
 MIP-1 alpha
 Prostatic AcidPhosphatase
 Testosterone

C) Analytes

Apolipoprotein B
 Apolipoprotein A2
 Apolipoprotein C1
 Cancer Antigen 125
 Calbindin
 CD5L
 EGF -R
 FSH (Follicle Stimulation Hormone)
 IgM
 IL-6 Receptor
 IL-11
 IL-17
 KIM-1
 MCP - 2
 MMP-2
 PYY
 Thyroid stimulating hormone
 Transferrin
 Vitronectin
 Thrombopoietin